

## SECTION II – GRAIN GRADING

### What is grain grading?

- Participants will be given 8 written scenarios of possible grain samples to be graded according to USDA standards. These descriptions will give grain type, weight, moisture, damages, foreign material, and other important information that is normally considered for the grain grading procedure.

### What do participants need to provide?

- Participants will need the following tools for this portion of the contest:
  - Clipboard
  - Pencil(s)
  - A non-scientific calculator

### What are participants provided with by contest officials?

- Grain grading problems
- Corn, Soybean and Wheat Charts (page 70-71)
- Official grain grading answer sheet (page 68-69)
- **Junior participants:** Grain grading handbook (pages 49-64)

### Rules for this portion of the contest

- This portion of the contest will have the following 8 grain samples:
  - 4 corn samples
  - 2 soybean samples
  - 2 wheat samples
- Participants will have 60 minutes to complete all 8 samples.
- Each sample is worth 25 points.
- Participants will grade samples based on the official grain grading handbook. This book will be provided to junior participants to use during this portion of the contest. Senior participants are not permitted a copy of the book during the contest at any time. (pages 49-64).

### How to prepare students for this portion of the contest

- Students should know all rules pertaining to grain grading found in the grain grading handbook. (pages 15-32).

For statistical information and other useful information regarding grain grading, see page 41.

## How to Grain Grade

- Proper spelling and capitalization is required.
  - The first letter of every word should be capitalized.
- See individual sections for rules on recording percentages and weights.

### How do you fill out the answer sheet?

Grade (required)	Class (required)	Special Grade Factors			Grade factors line				
GRADE AND KIND									
U.S. No. 1		Yellow Corn Infested							
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
56.0 lbs	13.4 %	0.2 %	1.7 %	0.1%	%	1.5 %	%	%	
Sample #1	Test Weight (required)	Moisture (required)	REMARKS						
							Other remarks and required information		

- Grade and Kind
  - Required for all samples
  - Should be stated as “U.S. No.” with designated number
  - Number is determined by charts after heat damage, damage kernels (total), special factors, etc. is calculated (charts on page 70-72)
  - If grade is sample grade, participants should strike out “No.” and write in “Sample Grade”

GRADE AND KIND									
U.S. <del>No.</del> Sample Grade		Yellow Corn							
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
56.0 lbs	13.4 %	%	%	%	%	%	%	%	
Sample #1	REMARKS Stones								

- Grain Class
  - Required for all samples
  - Varies for corn and soybeans. Wheat will **always** be Soft Red Winter Wheat.
  - The type of grain **MUST** be stated (ex. Yellow Corn, Mixed Soybeans)
- Test Weight and Moisture
  - Required for all samples, no matter the grade designation
  - Both numbers will be given in the sample
  - These numbers should be rounded to the nearest tenth for all samples

- Special grade factors
  - Varies for corn, soybeans, and wheat
  - These factors are listed in detail in the book
  - These factors normally do not affect the grade
  - If more than one special factor, list them in alphabetical order AFTER the grain class
  - In wheat, when dockage is present, it will always be listed last

GRADE AND KIND								
U.S. No. 3 Soft Red Winter Wheat, Ergoty, Infested, Dockage 6.2%								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*56.0 lbs	*14.1 %	%	%	%	%	%	%	%
Sample #7				REMARKS				

- Grade Factors Line
  - Fill out only those factors that contribute to the designation of the grade
  - Other boxes should be left blank
  - Not all boxes are used for all grain types (ex. In corn, the box “foreign material” will never be used)
  - *Note: Most answer keys to grain grading will list ALL factors and will place an asterisk or bold those answers required by the participant*

GRADE AND KIND								
U.S. No. 3 Soft Red Winter Wheat, Ergoty, Infested, Dockage 6.2%								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*56.0 lbs	*14.1 %	%	6.4 %	%	%	%	%	6.4 %
Sample #7				REMARKS				

- Other Remarks
  - This box is used for 3 types of information: percentage of mixed classes, sample grade factors, and any factor for which there is no other box (ex. Soybeans of Other Colors).
  - For mixed samples, sample type should be listed first and listed from greatest to least
  - All others should be listed in alphabetical order
  - See manual for specific information that should be in this box

GRADE AND KIND								
U.S. No. Sample Grade Mixed Corn								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*56.0 lbs	*12.1 %	%	%	%	%	%	%	%
Sample #3				REMARKS Yellow Corn 85.0% White Corn 15.0% Cockleburrs				

### How do you use the charts?

Minimum test weight per bushel

- Test weight listed for grades is the **minimum** amount a grain can weigh
- Example: Corn that is U.S. No. 1 can weigh 56 lbs. or more, but it cannot weigh less.
- Example: Corn that weighs 55.9 lbs. it is U.S. No. 2.
- **Note**
  - All wheat samples will be Soft Red Winter Wheat, therefore, participants will NEVER use the test weight category of “Hard Red Spring Wheat or White Club Wheat”
  - Soybean test weight is no longer a factor in determining grades

Other categories

- Maximum limits are given for all other factors. This means a sample cannot contain more than a given maximum percentage of a given factor.
- Example: Corn that is U.S. No. 1 cannot have more than 2.0% Broken Corn and Foreign Material
- Example: Corn that has 2.1% Broken Corn and Foreign Material must be U.S. No. 2.

### What is the final grade of your grain?

The factor resulting in the lowest grade determines the grade for the lot of grain. **Samples that are U.S. No. 1 do not require factors listed except the test weight, moisture, and special factors that do not contribute to the grade designation.**

Example: yellow soybean sample

Test Weight (lbs/bu)	51.0	<b>No effect</b>
Foreign Material	0.8 %	<b>U.S. No. 1</b>
Moisture	13.2%	<b>No effect</b>
Splits	18.7%	<b>U.S. No. 2</b>
Green Garlic bulbs	6	<b>Garlicky</b>

This sample would be U.S. No. 2 and will be recorded as follows:

GRADE AND KIND								
U.S. No. 2 Yellow Soybeans, Garlicky								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*51.0 lbs	*13.2 %	%	%	%	18.7 %	%	%	%
Sample #6				REMARKS				

If, for any reason, the sample fails to make any of the numerical grades, the sample is graded “U.S. Sample Grade.” Factors for which any sample is automatically graded as Sample Grade are listed in the manual and below each chart.

Example: Mixed Corn (Yellow Corn 80.0%, White Corn 20.0%)  
 Test Weight (lbs/bu) 55.0 U.S. No. 2  
 Foreign Material 5.1 % U.S. Sample  
 Moisture 13.2% No effect  
 Damaged Kernels (Total) 1.0% U.S. No. 1

This sample would be U.S. Sample Grade and will be recorded as follows:

GRADE AND KIND									
U.S. <del>No.</del> Sample Grade		Mixed Corn							
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
55.0 lbs	13.2 %	%	1.0%	5.1%	%	%	%	%	
Sample #2	<i>Since this makes it a U.S. No. 1, it is not required on the answer sheet; however, if it is included and is the correct number, no points will be deducted</i>			REMARKS Yellow Corn 80.0% White Corn 20.0%					

### How is the answer sheet scored?

- *The Official Grain Standards of the United States*, published by the U.S. Department of Agriculture, Federal Grain Inspection Service, will be the official manual for completing the answer sheet which resembles an official grain grading certificate. If, however, the rules of the grading change during the five year coverage of this document, the rules spelled out within will have presidency.
- Some customers request that information in addition to grading factors be recorded on the grain certificate. If extra information (other than factors which affect grade) is recorded on the answer sheet, the information **MUST** be correct. **If the additional information is not correct, then 1 point is deducted for each incorrect factor box.**
  - Example: If a participant incorrectly records Damage Kernels (Total) Information and that information is not a grading factor, the participant would lose 1 of their 12 points possible for that line.
- Incorrectly written percentages will result in an incorrect answer.
  - Numbers less than 1 should be written to include a zero before the decimal
    - Ex. .8% should be recorded at 0.8%
    - Contestant will lose all points if incorrect
- Extraneous marks, such as zeros or dashes, put in factor boxes will be counted as wrong answers. This is for the benefit of graders in that they do not have to sift through an over amount of extra information. Will be counted off as minus 1.
- The sample will be scored on the basis of grade line, grading factors line, and remarks section for a total of 25 points per sample (200 points total).
- Negative score for any line will be scored as zero points

### **How to score corn and soybean problems**

- Grade Line-10 points total
  - Grade-6 points
    - Incorrect by 1 grade: 2 points off
    - Incorrect by 2 grades: 4 points off
    - Incorrect by 3 or more grades: 6 points off
  - Correct Class- 4 points
    - Yellow, White, or Mixed Corn
    - Yellow or Mixed Soybeans
  - Special Factors- 3 points
    - Specific for each grain type
- Grading Factors Line-12 points total
  - All correct: 12 points
  - 2 factors required with 1 correct: 6 points off
  - 3 factors required with 2 correct: 4 points off
  - 3 factors required with 1 correct: 8 points off
  - 4 factors required with 3 correct: 3 points off
  - 4 factors required with 2 correct: 6 points off
  - 4 factors required with 1 correct: 9 points off
  - No factors correct: 12 points off
- Remarks section- 3 points total
  - Specific for each grain type

### **How to score wheat problems**

- Grade Line-10 points total
  - Grade-6 points
    - Incorrect by 1 grade: 2 points off
    - Incorrect by 2 grades: 4 points off
    - Incorrect by 3 or more grades: 6 points off
  - Correct Class- 1 point
    - Soft Red Winter Wheat
  - Special Factors- 3 points
    - Specific for each grain type
  - Dockage correctly recorded- 3 points
- Grading Factors Line-12 points total
  - See “how to grade corn and soybean problems”
- Remarks section- 3 points total
  - Specific for each grain type

**What is corn?**

Corn is defined as any grain which consists of 50 percent or more of whole kernels of shelled dent corn and/or flint corn. It may not contain more than 10 percent of other grains for which grading standards have been established. If it does not meet these standards, the lot is considered mixed grain. However, in this event there will be no mixed grain so any grain other than dent or flint corn is foreign material. **Popcorn, sweet corn, and blue corn in corn grading are foreign material.**

Class and damaged kernels are determined **after** the removal of foreign material. All percentages shall be determined on the grain as a whole.

**Corn Grain Class**

There are three possible classes of corn, Yellow Corn, White Corn and Mixed Corn. Percentage of corn classes should be rounded to the nearest tenth.

**Yellow Corn** - Yellow-kernelled corn that does not contain **more than 5 percent** corn of other colors. In other words, 95.0% or more Yellow Corn. Yellow kernels with red streaks **covering less than 50.0%** of the kernel are considered yellow corn.

**White Corn** - White-kernelled corn that does not contain **more than 2 percent** corn of other colors. In other words, 98.0% or more White Corn. White corn with a slight tinge of pink is white corn.

**Mixed Corn** - Corn that does not meet the color requirements of white or yellow corn. When completing the grain inspection certificate (answer sheet) record the percent of each corn (class) from greatest to least in percentage to nearest tenth within the "Remarks" section. If more than 50.0% of the kernel is red streaked, then the corn is considered Red Corn. Students should write **"Red Corn"** in the remarks section if it is a determining factor for corn grain class.

GRADE AND KIND									
U.S. No. 1 <b>Yellow Corn</b>									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*13.4 %	%	%	%	%	%	%	%	%
Sample #1					REMARKS				

GRADE AND KIND									
U.S. No. 1 <b>Mixed Corn</b>									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*11.2 %	%	%	%	%	%	%	%	%
Sample #2					REMARKS				
					Yellow Corn 85.0% White Corn 15.0%				

### Moisture

Moisture is not a grading factor in commercial grain; nevertheless, a loss of quality in stored corn hinges largely on the amount of moisture present in the grain. Moisture is an important factor in most discount schedules.

Moisture is recorded to the nearest tenth of a percent.

- Example: **16.27% is recorded as 16.3%**

### Test Weight

Test weight is the amount of weight the grain must have to make up a bushel. Good quality corn of low moisture content can be expected to have a good test weight.

Test weight is recorded to the nearest tenth.

- Example: **52.34 lbs/bu is recorded as 52.3 lbs/bu**

### Broken Corn and Foreign Material

Broken corn and foreign material is normally determined by the use of a sieve; broken corn and all matter other than corn that pass through a sieve having round openings 12/64th of an inch in diameter, and all matter other than corn that remain on the sieve after screening are included in this factor.

Examples of foreign material

- Sweet corn
- Popcorn
- Blue corn
- Soybeans not passing through the sieve
- Grains or weeds dropping through the sieve
- Rodent excreta and stones (cinders are stones)
  - *Note:* If the total weight of stones in a 1-1/4 quarts (1,000 gm.) sample **exceeds** 0.1 percent of the sample weight the sample must be graded "U.S. Sample Grade". When applicable, record "**Stones**" in the Remarks section of the certificate.

GRADE AND KIND								
U.S. No. <b>Sample Grade</b> <b>Yellow Corn</b>								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*56.0 lbs	*14.1 %	%	%	%	%	%	%	%
Sample #3				REMARKS				
				Stones				

**Do not enter this percentage in the foreign material column on the answer sheet.** There is a column for broken corn and foreign material.

Broken corn and foreign material is recorded to nearest tenth of a percent.



## Heat Damaged Kernels

Heat damaged corn is **severely** discolored (brown to black) either from external heating, such as improper drying, or from heating as a result of excessive moisture in storage and spoilage. Kernels in this category will be included in heat damage **AND** damaged kernels (total).

Slightly damaged corn shows some discoloration (light to dark tan) and therefore is not as severely damaged. The two are **not** added together to determine heat damaged kernels. Slight damaged corn will only be included in damaged kernels (total).

Record to the nearest tenth of a percent.

## Damaged Kernels (Total)

Damaged kernels (total) includes **all** types of damage found in corn. Darkening of the germ is one of the first indicators of corn declining in quality or that the amount of damage is increasing.

Examples of damaged kernels

- mold damage
- heat damage
- sprout damage
- frost damage
- badly ground-damaged
- badly weather-damaged
- insect damage (not chewed)
- kernels that have become slightly discolored from heat

Note that the percent of heat damage is added to other types of damage to obtain the percent of Damaged Kernels (Total).

Record to the nearest tenth of a percent.

## Musty, Sour, or Heating

A sample in any of these conditions is "U.S. Sample Grade."

Musty- Musty, ground, or moldy odor

Sour- Sour, fermenting, or pigpen odor

Heating- Corn developing a high temperature from excessive respiration. Corn will usually have a musty or sour odor. For this contest, samples that are affected by heating will state "corn affected by high temperature" OR "heating."

Record the applicable words in the "Remarks" section of the grading certificate.

GRADE AND KIND								
U.S. No. Sample Grade Yellow Corn								
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)
*56.0 lbs	*14.1 %	%	%	%	%	%	%	%
Sample #3					REMARKS Heating Musty Sour			

**Commercially Objectionable Foreign Odor**

If the corn carries an odor which does not normally occur in grain and which, for this reason, would render the corn unfit for its normal commercial use, then it is graded "U.S. Sample Grade."

This includes animal hides, decaying animal or vegetable parts, fertilizer, skunk, smoke, strong weed, oil, etc.

Record the words "Commercially Objectionable Foreign Odor" in the "Remarks" section of the certificate.

GRADE AND KIND									
U.S. No. <b>Sample Grade</b> <b>Yellow Corn</b>									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*14.1 %	%	%	%	%	%	%	%	
Sample #3				REMARKS Commercially Objectionable Foreign Odor					

**Distinctly Low Quality**

The Federal Grain Inspection Service reserves the use of this term to describe corn when it is obviously of inferior quality and the existing grading factors or guidelines do not accurately reflect the inferior condition.

When a sampler is collecting corn from a rail car, he/she can notice whether the grain also includes two or more large stones, pieces of glass, pieces of concrete, sticks of lumber, or scrap metal or debris which are visible to the sampler but are too large to enter the sampling device, such as a grain probe.

This grading factor **should not** be confused with the other conditions which can also cause corn to be "Sample Grade," such as animal filth, cockleburs, crotalaria seed, etc. Record the words Distinctly Low Quality (Reason).

GRADE AND KIND									
U.S. No. <b>Sample Grade</b> <b>Yellow Corn</b>									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*14.1 %	%	%	%	%	%	%	%	
Sample #3				REMARKS Distinctly Low Quality (Metal Scraps)					

## Sample Grade Factors

There is a list of factors that does not meet U.S. number standards and make the sample "Sample Grade" listed on the bottom of the grading chart. These include animal filth, cockleburs, crotalaria seed. If any of these are reported in the sample, participants should grade the sample "Sample Grade" and write the word in the Remarks box.

How to record in the "Remarks" box (see chart for details)

- Stones
- Glass
- Crotalaria
- Castor Beans
- Unknown Foreign Substance
- Commonly Recognized Harmful or Toxic Substance
- Cockleburs
- Animal Filth (Includes bird droppings and rodent pellets)
- Heating
- Distinctly Low Quality (Reason)

## Special Factors, Special Grade Requirements, Special Grade Designations

Special grades are conditions which should be noted but **do not** affect the numerical grade. They should be listed in alphabetical order with commas in between.

### Flint

Corn of any class which consists of **95 percent or more** of flint corn; flint corn is graded and designated according to the grade requirements of the standards applicable to such corn if it were not flint, and the word "Flint" is added to and made a part of the grade designation, immediately following the class name.

Flint corn is a different subspecies of corn with hard starch rather than soft starch as in dent corn.

GRADE AND KIND									
U.S. No. 1 Yellow Corn, Flint, Infested									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*12.3 %	%	%	%	%	%	%	%	
Sample #1					REMARKS				

### Flint and Dent

Corn of any class which consists of a mixture of flint and dent corn containing **more than 5 percent but less than 95 percent of flint corn**. The words "Flint and Dent" and the percentage of flint corn rounded to the tenth are added to and made a part of the grade designation immediately following the special grade.

GRADE AND KIND									
U.S. No. 1 Yellow Corn, Flint and Dent, Flint Corn 20.0%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*12.3 %	%	%	%	%	%	%	%	
Sample #1					REMARKS				

**Infested**

Any corn sample 1-1/4 quarts or 1000 g that contains one of the following:

- 2 or more live weevils
- 1 live weevil and 5 or more other live insects injurious to stored grain
- 10 or more other live insects injurious to stored grain

Infested is the condition of live weevils or grain-damaging insects in the grain.

Infested corn is graded and designated according to the grade requirements of the standards applicable to such corn if it was not infested. The word "Infested" is added to and made a part of the grade designation.

**Waxy Corn**

Corn that consists of **95% or more waxy corn**. When applicable, the word "Waxy" will be last within the grade designation.

GRADE AND KIND									
U.S. No. 1 Yellow Corn, Flint, Infested, Waxy									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*13.7 %	%	%	%	%	%	%	%	
Sample #2					REMARKS				

GRADE AND KIND									
U.S. No. 1 Yellow Corn, Flint and Dent, Flint Corn 15.0%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*13.7 %	%	%	%	%	%	%	%	
Sample #2					REMARKS				

## Soybean Grading

### What are soybeans?

Soybeans are any grain that consists of 50 percent or more of whole or broken soybeans which will not pass readily through an 8/64 sieve and not more than 10 percent of other grains for which grading standards have been established.

Class, splits, and damaged kernels are determined after foreign material is removed.

### Soybean Grain Class

There are two possible classes of soybeans, Yellow Soybeans or Mixed Soybeans. Percentage of soybean classes should be rounded to the **nearest tenth**.

**Yellow Soybeans** - Soybeans that have a yellow seed coat and are yellow in cross-section. Sample does not contain more than 10 percent of other colors.

**Mixed Soybeans** - Any mixture of soybeans that does not meet the requirements of yellow soybeans (See Soybeans of Other Colors). When completing the grain inspection certificate, record the percent of each color of soybeans from greatest to least to the nearest tenth within the Remarks section.

**Soybeans of Other Colors**- These colors serve as a **grading factor** in yellow soybeans. When soybeans of other colors (black, brown, and bi-colored) **occur in quantities of 10 percent or less**, the percentage is a factor in determining the grade of yellow soybeans. **When other colors exceed 10 percent, the sample is then classified as Mixed Soybeans** (see above). Soybeans of other colors is listed in the Remarks section.

### Example

Yellow Soybeans	85.0%	Mixed Soybeans
Bi-color Soybeans	15.0%	
Yellow Soybeans	95.0%	Yellow Soybeans
Bi-color Soybeans	5.0%	Soybeans Of Other Colors

### Test Weight

Test weight is the amount of weight the grain must have to make up a bushel. Good quality seed of low moisture content can be expected to have a good test weight. **Test weight has no effect on the grade of soybean samples.**

Record test weight rounded to the nearest tenth.

### Moisture

The moisture content of soybean seed is extremely important but it is no longer used as a grading factor. Loss of quality of stored seed hinges largely on the amount of moisture present in the sample. Moisture is an important factor in most discount schedules.

Record moisture to the nearest tenth of a percent.

## Splits

Any soybean having **more than** 1/4 of the seed missing is considered a split. Splits are determined on a portion of approximately 125 grams after the removal of all foreign material.

This factor includes only **sound splits** - those free from damage.

Damaged splits are **only** recorded in Damaged Kernels (Total).

Splits are recorded to the nearest tenth of a percent.

## Heat Damage

Heat damaged soybeans are **severely** discolored (black or dark brown) either from external heating, such as improper drying, or from heating as a result of excess moisture and spoiling. Almost all heat damage is the result of storing grain too wet. Soybeans in this category will be included in heat damage **AND** Damaged Kernels (Total).

Slightly damaged soybeans show some discoloration (light to dark tan) and therefore are not as severely damaged. The two are **not** added together to determine heat damaged kernels. Slight damaged soybeans will only be included in Damaged Kernels (Total).

Record heat-damaged kernels to the nearest tenth of a percent.

## Damaged Kernels (Total)

This factor includes **all** types of damage found in whole and pieces of soybeans.

### Examples of Damaged Kernels

- heat damage
- sprout damage
- frost damage
- immature seed
- ground-damage
- mold damage
- insect damage (not chewed)
- kernels that have become slightly discolored from heat
- heat damage
- stink bug stung kernels\*\*

\*\*Stink bug stung kernels are considered damaged kernels at the rate of 1/4 of actual total percentage of stung kernels

Example: 12 grams is considered as 3 grams of damage

Record Damage Kernels (Total) to the nearest tenth of a percent.

## Foreign Material

Foreign material is normally determined by the use of a sieve and separated into coarse and fine foreign material.

Coarse foreign material includes material that does not pass through an 8/64 inch sieve and made on 1000 grams.

Fine foreign material includes material that passes through an 8/64 inch sieve and material and pieces of soybean that remains on top of the sieve after sieving. This test is made on 125 grams.

Examples of foreign material

- corn
- cockleburrs
- sticks
- stalks
- rodent excreta
- stones
- other grains

Record to the nearest tenth of a percent.

## Distinctly Low Quality

The Federal Grain Inspection Service reserves the use of this term to describe soybeans when it is obviously of inferior quality and the existing grading factors or guidelines do not accurately reflect the inferior condition.

When a sampler is collecting soybeans from a rail car, he/she can notice whether the grain also includes two or more large stones, pieces of glass, pieces of concrete, sticks of lumber, or scrap metal or debris which are visible to the sampler but are too large to enter the sampling device, such as a grain probe.

This grading factor **should not** be confused with the other conditions which can also cause soybeans to be "Sample Grade", such as animal filth, cockleburrs, crotalaria seed, etc. Record the words Distinctly Low Quality (Reason).

*(See corn for example of how to record).*

## Musty, Sour or Heating

A sample in any of these conditions is "U.S. Sample Grade".

Musty- Musty, ground, or moldy odor

Sour- Sour, fermenting, or pigpen odor

Heating- Soybeans developing a high temperature from excessive respiration. Soybeans will usually have a musty or sour odor. For this contest, samples that are affected by heating will state "Soybeans affected by high temperature" OR "heating."

Record the applicable words in the "Remarks" section of the grading certificate.

*(See corn for example of how to record)*

### Commercially Objectionable Foreign Odor

If the soybeans carry an odor which does not normally occur in grain and which, for this reason, would render the soybeans unfit for its normal commercial use, then it is graded "U.S. Sample Grade."

This includes animal hides, decaying animal or vegetable parts, fertilizer, skunk, smoke, strong weed, oil, etc.

Record the words "Commercially Objectionable Foreign Odor" in the "Remarks" section of the certificate.

(See corn for example of how to record)

### Sample Grade Factors

There is a list of factors that automatically make the sample "Sample Grade" listed on the bottom of the grading chart. These include animal filth, cocklebur, crotalaria seed. If any of these are reported in the sample, participants should grade the sample "Sample Grade" and write the word in the "Remarks" box.

How to record in the "Remarks" box (see chart for details)

- Stones
- Glass
- Crotalaria
- Castor Beans
- Unknown Foreign Substance
- Commonly Recognized Harmful or Toxic Substance
- Cocklebur
- Animal Filth (Includes bird droppings and rodent pellets)
- Heating
- Distinctly Low Quality (Reason)

### Cumulative Total

If a cumulative total of 11 or more sample grade factors are found, the sample is graded "U.S. Sample Grade."

These factors include any combination of animal filth, castor beans, crotalaria seeds, glass, stones, and any unknown foreign substance.

When applicable, record "**Cumulative Total**" in the "Remarks" section. You are **not required** to list the factors that cumulate to this total.

GRADE AND KIND									
U.S. No. Sample Grade Yellow Soybeans									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*16.0 %	%	6.0%	%	%	%	%	%	
Sample #6					REMARKS Cumulative Total Soybeans of Other Colors 6.5%				



## Special Factors, Special Grade Requirements, Special Grade Designations

Special grades are conditions which should be noted but **do not** affect the numerical grade. They should be listed in alphabetical order with commas in between.

### Garlicky

Specific types of garlic bulbs found in the sample may be considered a special factor.

Green bulbs- husk is still intact, contains **three or more** green bulblets in 1000 grams

Dry bulbs- husk is dry or missing, **3 dry bulbs= 1 green bulb**

The word "Garlicky" is added to and made a part of the grade designation. A garlic odor is not a basis for "Garlicky."

### Infested

Any soybean sample that contains one of the following:

- two or more live weevils
- one live weevil and 5 or more other live insects injurious to stored grain
- 10 or more other live insects injurious to stored grain

Infested is the condition of live weevils or grain-damaging insects in the grain.

Infested soybeans are graded and designated according to the grade requirements of the standards applicable to such soybeans if it was not infested. The word "Infested" is added to and made a part of the grade designation.

### Purple Mottled or Stained

Soybeans that are discolored with pink or purple seed coats, dirt or dirt-like substance, or pokeberry stains, as determined on a portion of 400 grams with the use of an FGIS Interpretive Line Print. The Interpretive Line Prints are help tools for inspectors to judge whether or not a kernel is damage or not.

Samples with this condition will state "Purple Mottled or Stained" as the description.

The words "Purple Mottled or Stained" are added to and made part of the grade designation.

GRADE AND KIND									
U.S. No. 1 Mixed Soybeans, Garlicky, Infested, Purple Mottled or Stained									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*56.0 lbs	*13.7 %	%	0.3 %	%	%	%	%	%	
Sample #6					REMARKS				
					Yellow Soybeans 85.6% Black Soybeans 10.0% Bi-Color Soybeans 4.4%				

## Wheat Grading

### What is Wheat?

Wheat is any grain of common wheat, club wheat, and durum wheat, which before the removal of dockage, consists of 50 percent or more of these wheats and not more than 10 percent of other grains for which standards have been established and which, after the removal of dockage, contains 50 percent or more of whole kernels of one or more of these wheats.

Wheat is divided into the following seven classes: Hard Red Spring Wheat, Durum Wheat, Soft Red Winter Wheat, Hard Red Winter Wheat, Hard White Wheat, Soft White Wheat, Unclassed Wheat, and Mixed Wheat.

**Soft Red Winter Wheat is the only wheat class that will be graded in this event**, and the following discussion will pertain **only** to Soft Red Winter Wheat. The class Soft Red Winter Wheat includes all varieties of Soft Red Winter Wheat. There are no subclasses in this class.

### Basis of Determination

Each determination of dockage, moisture, temperature, odor, garlic, live weevils or other insects injurious to stored grain, and distinctly low quality completed on the grain as received when taken from an incoming truck, rail car, etc. All other "tests" are conducted after dockage has been removed.

### Test Weight

Test weight is the amount of weight the grain must have to make up a bushel. Good quality wheat of low moisture content can be expected to have a good test weight.

Record test weight rounded to the nearest tenth of a percent.

### Moisture

The moisture content of wheat seed is extremely important but it is no longer used as a grading factor. Loss of quality of stored seed hinges largely on the amount of moisture present in the sample. Moisture is an important factor in most discount schedules.

Moisture is recorded to the nearest tenth of a percent.

### Dockage

The word "dockage" means weed seed, weed stems, chaff, straw, grain other than wheat, sand, soil, and any other material other than wheat, that can be removed readily from the wheat by the use of appropriate sieves and cleaning devices. Also, the underdeveloped, shriveled, and small pieces of wheat kernels removed in properly separating the material other than wheat plus that which cannot be recovered by properly rescreening or re-cleaning is also a part of dockage.

Determination of dockage is made in the initial sieving. Shrunken and broken kernels and foreign material are determined after the dockage has been removed. Dockage is determined from a 1,000 gram sample. ***Participants will need to convert the number to a percentage if it is reported in grams.***

The percent dockage is rounded and reported to the nearest tenth percent. Always list dockage last of all of the special factors. See example on the special factors page.

### Foreign Material

Foreign material refers to all matter other than wheat, including stones, that is not separated from the wheat in the proper removal of dockage.

Examples of foreign material

- corn
- cocklebur
- sticks
- stalks
- rodent excreta
- stones
- other grains
- ergot wheat

Record to the nearest tenth of a percent.

### Contrasting Classes

Contrasting classes in soft red winter wheat are durum wheat and unclassified wheat.

Soft red winter wheat flour is especially suited for cake mixes while flour from durum wheat is required for pasta production. Thus, there is a "contrast" in use. Each wheat has its own "Contrasting Classes."

Record in Remarks area of certificate "Contrasting Classes" and state to the nearest tenth of a percent.

GRADE AND KIND									
U.S. No. 1 Soft Red Winter Wheat, Dockage 12.4%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*60.0 lbs	*13.3%	%	%	%	%	%	%	%	
Sample #8				REMARKS Contrasting Classes 0.8%					

### Wheat of Other Classes (Total)

This factor spotlights the presence of other wheats in a sample. Some mixtures may be of minor importance. For example, if a soft red winter wheat contained 8.0% hard red winter wheat, the flour from such a mixture might be acceptable, but not the most desirable for cake mixes when compared to flour from 100.0% soft red winter wheat.

Wheat of Other Classes (Total) also includes percent of Contrasting Classes.

Record "Wheat of Other Classes (Total)" and state to the nearest tenth of a percent in the "Remarks" section.

GRADE AND KIND									
U.S. No. 1 Soft Red Winter Wheat, Dockage 12.4%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*60.0 lbs	*13.3%	%	%	%	%	%	%	%	
Sample #8				REMARKS Wheat of Other Classes (Total) 1.2%					

## Other Grains

Other grains as used in this discussion are:

- rye
- oats
- corn
- grain sorghum
- barley
- flax
- emmer
- spelt
- einkorn
- polish wheat
- poulard wheat
- cultivated buckwheat
- soybeans

These grains are also considered foreign material, even when damaged.

## Heat Damage

Heat damage in wheat is **severely** discolored (black or dark brown) kernels and pieces of kernels of **wheat and other grains** caused either from external heating, such as improper drying, or from heating as a result of excess moisture and spoiling. Almost all heat damage is the result of storing grain too wet. All grains listed in Other Grains above that are severely heat damaged **in Wheat** will be included in heat damage, damaged kernels (total), **and** Foreign Material.

Slightly damaged wheat and other grains show some discoloration (light to dark tan) and therefore are not as severely damaged. The two are **not** added together to determine heat damaged kernels. Slight damaged wheat and other grains will only be included in damaged kernels (total).

Record heat-damaged kernels to the nearest tenth of a percent.

## Insect Damaged Wheat Kernels

Wheat is determined to be "U.S. Sample Grade" when 32 or more insect damaged kernels per 100 grams are found. This is up to a 3 stage process. For simplicity in this event, the number of kernels per 100 gram will be given.

Do not confuse insect chewed with insect damage. Insect damage includes drill holes, pin holes, bored, tunneling and webbing in description.

When applicable, include in the "Remarks" section of the certificate "**Insect Damaged Kernels.**"

## Damaged Kernels (Total)

This factor includes all types of damage found in wheat. It is very inclusive in that kernels and pieces of kernels of **wheat plus other grains (Ex. Sprout-damaged Oats)** are also included.

Examples of Damaged Kernels

- heat-damage
- sprout damage
- frost damage
- badly ground-damage
- badly weather-damage
- mold damage
- insect damage (not chewed)
- disease or otherwise materially damaged

Damaged Kernels (Total) is recorded to the nearest tenth of a percent.

### **Shrunken and Broken Kernels**

These are kernels and pieces of kernels of wheat and other matter that will pass readily through a .064 x 3/8 inch oblong hole sieve after the dockage has been removed.

Record to the nearest tenth of a percent.

### **Defects (Total)**

This factor is determined by adding the percentages of Damaged Kernels (Total), Foreign Material, and Shrunken and Broken Kernels.

### **Distinctly Low Quality**

The Federal Grain Inspection Service reserves the use of this term to describe wheat when it is obviously of inferior quality and the existing grading factors or guidelines do not accurately reflect the inferior condition.

When a sampler is collecting wheat from a rail car, he/she can notice whether the grain also includes two or more large stones, pieces of glass, pieces of concrete, sticks of lumber, or scrap metal or debris which are visible to the sampler but are too large to enter the sampling device, such as a grain probe.

This grading factor **should not** be confused with the other conditions which can also cause wheat to be "Sample Grade," such as animal filth, cocklebur, crotalaria seed, etc.

Record the words Distinctly Low Quality (Reason).

### **Musty, Sour or Heating**

A sample in any of these conditions is "U.S. Sample Grade."

Musty- Musty, ground, or moldy odor

Sour- Sour, fermenting, or pigpen odor

Heating- Wheat developing a high temperature from excessive respiration. Wheat will usually have a musty or sour odor. For this contest, samples that are affected by heating will state "Wheat affected by high temperature" OR "heating."

Record the applicable words in the "Remarks" section of the grading certificate.

### **Commercially Objectionable Foreign Odor**

If the wheat carries an odor which does not normally occur in grain and which, for this reason, would render the wheat unfit for its normal commercial use, then it is graded "U.S. Sample Grade."

This includes animal hides, decaying animal or vegetable parts, fertilizer, skunk, smoke, strong weed, oil, etc. **This does not include smutty or garlicky odor.**

Record the words "Commercially Objectionable Foreign Odor" in the "Remarks" section of the certificate.

### Sample Grade Factors

There is a list of factors that automatically make the sample "Sample Grade" listed on the bottom of the grading chart. These include animal filth, cocklebur, crotalaria seed. If any of these are reported in the sample, participants should grade the sample "Sample Grade" and write the word in the "Remarks" box.

How to record in the "Remarks" box (see chart for details)

- Stones
- Glass
- Crotalaria
- Castor Beans
- Unknown Foreign Substance
- Commonly Recognized Harmful or Toxic Substance
- Cockleburs
- Animal Filth (includes bird droppings and rodent pellets)
- Heating
- Distinctly Low Quality (Reason)

### Cumulative Total

If a cumulative total of 5 or more sample grade factors (e.g. 3 stones + 1 animal filth + 1 unknown = 5 or more sample grade factors) are found, the sample is graded "U.S. Sample Grade." Record "**Cumulative Total**" in the "Remarks" section.

GRADE AND KIND									
U.S. No. Sample Grade Soft Red Winter Wheat, Dockage 12.4%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*60.0 lbs	*13.3 %	%	%	%	%	%	%	%	
Sample #8				REMARKS					
				Cumulative Total					

## **Special Factors, Special Grade Requirements, Special Grade Designations**

Special grades are conditions which should be noted but **do not** affect the numerical grade. All special grades should be listed in alphabetical order with commas in between.

### **Ergoty**

Wheat that contains **more than 0.05 percent** per 1000 grams ergot is considered Ergoty.

The word "Ergoty" is added to and made part of the grade designation.

Note that **ergot also fits the definition of foreign material** in wheat and must be included as such.

### **Garlicky**

Specific types of garlic bulbs found in the sample may be considered a special factor.

Green bulbs- husk is still intact, contains **three or more** green bulblets in 1000 grams

Dry bulbs- husk is dry or missing, **3 dry bulbs= 1 green bulb**

The word "Garlicky" is added to and made a part of the grade designation. A garlic odor is not a basis for "Garlicky."

### **Infested**

Any wheat sample that contains one of the following:

- two or more live weevils
- one live weevil and 1 or more other live insects injurious to stored grain
- 2 or more other live insects injurious to stored grain

Infested is the condition of live weevils or grain-damaging insects in the grain.

Infested wheat is graded and designated according to the grade requirements of the standards applicable to such wheat if it was not infested. The word "Infested" is added to and made a part of the grade designation.

### **Smutty**

There are two special grades of smutty wheat -- Light Smutty and Smutty.

- Light Smutty - Applies to wheat with a smutty odor, or when wheat contains 6-30 smut balls in 250 grams of grain; the term "Light Smutty" is added to and made part of the grade designation.
- Smutty - Applies to wheat that contains 31 or more smut balls per 250 gram sample; the word "Smutty" is added to and made part of the grade designation.

### **Treated Wheat**

Treatments of wheat include:

- Scoured
- Washed
- Limed
- Sulphured

Wheat that is treated will have the words "Treated (treatment(s))" as part of the grade



designation.

**Wheat Special Factors Examples**

GRADE AND KIND									
U.S. No. 1 Soft Red Winter Wheat, Garlicky, Light Smutty, Treated (Limed), Dockage 12.4%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*60.0 lbs	*13.3 %	%	%	%	%	%	%	%	
Sample #8				REMARKS					

GRADE AND KIND									
U.S. No. 1 Soft Red Winter Wheat, Ergoty, Smutty, Treated (Washed), Dockage 12.4%									
TEST WEIGHT PER BUSHEL	MOISTURE	HEAT DAMAGED KERNELS	DAMAGED KERNELS (TOTAL)	FOREIGN MATERIAL	SPLITS	BROKEN CORN AND FOREIGN MATERIAL	SHRUNKEN AND BROKEN KERNELS	DEFECTS (TOTAL)	
*60.0 lbs	*13.3 %	%	%	%	%	%	%	%	
Sample #8				REMARKS					

**See page 43 for an example of the grain grading problem set.**

**Past grain grading problems from invitational contests, area, and state can be found on the agronomy website (see reference page for information).**